

# Nonlinear Preconditioning in PETSc

Matthew Knepley ∈ PETSc Team

Computation Institute  
University of Chicago

Department of Molecular Biology and Physiology  
Rush University Medical Center

Earth Science & Engineering Seminar  
Imperial College London, UK   March 18, 2014



# The PETSc Team



Bill Gropp



Barry Smith



Satish Balay



Jed Brown




Matt Knepley



Lisandro Dalcin



Hong Zhang



Mark Adams



Toby Isaac

# Outline

1 Algorithmics

2 Experiments

# Abstract System

Our prototypical nonlinear equation is:

$$\mathcal{F}(\vec{x}) = \vec{b} \quad (1)$$

and we define the residual as

$$\vec{r}(\vec{x}) = \mathcal{F}(\vec{x}) - \vec{b} \quad (2)$$

# Abstract System

Our prototypical nonlinear equation is:

$$\mathcal{F}(\vec{x}) = \vec{b} \quad (1)$$

and we define the (linear) residual as

$$\vec{r}(\vec{x}) = A\vec{x} - \vec{b} \quad (3)$$

# Linear Left Preconditioning

The modified equation becomes

$$P^{-1} (\vec{A}\vec{x} - \vec{b}) = 0 \quad (4)$$

# Linear Left Preconditioning

The modified defect correction equation becomes

$$P^{-1} (A\vec{x}_i - \vec{b}) = \vec{x}_{i+1} - \vec{x}_i \quad (5)$$

# Nonlinear Additions

Unlike the linear case, we must define

- the solution  $\vec{x}$

AND

- the residual  $\vec{r}$

in both the inner and outer solvers.

# Additive Combination

The linear iteration

$$\vec{x}_{i+1} = \vec{x}_i - (\alpha P^{-1} + \beta Q^{-1})(A\vec{x}_i - \vec{b}) \quad (6)$$

becomes the nonlinear iteration

# Additive Combination

The linear iteration

$$\vec{x}_{i+1} = \vec{x}_i - (\alpha P^{-1} + \beta Q^{-1}) \vec{r}_i \quad (7)$$

becomes the nonlinear iteration

# Additive Combination

The linear iteration

$$\vec{x}_{i+1} = \vec{x}_i - (\alpha P^{-1} + \beta Q^{-1}) \vec{r}_i \quad (7)$$

becomes the nonlinear iteration

$$\vec{x}_{i+1} = \vec{x}_i + \alpha(\mathcal{N}(\mathcal{F}, \vec{x}_i, \vec{b}) - \vec{x}_i) + \beta(\mathcal{M}(\mathcal{F}, \vec{x}_i, \vec{b}) - \vec{x}_i) \quad (8)$$

# Nonlinear Left Preconditioning

From the additive combination, we have

$$P^{-1}\vec{r} \implies \vec{x}_i - \mathcal{N}(\mathcal{F}, \vec{x}_i, \vec{b}) \quad (9)$$

so we define the preconditioning operation as

$$\vec{r}_L \equiv \vec{x} - \mathcal{N}(\mathcal{F}, \vec{x}, \vec{b}) \quad (10)$$

# Multiplicative Combination

The linear iteration

$$\vec{x}_{i+1} = \vec{x}_i - (P^{-1} + Q^{-1} - Q^{-1}AP^{-1})\vec{r}_i \quad (11)$$

becomes the nonlinear iteration

# Multiplicative Combination

The linear iteration

$$\vec{x}_{i+1/2} = \vec{x}_i - P^{-1} \vec{r}_i \quad (12)$$

$$\vec{x}_i = \vec{x}_{i+1/2} - Q^{-1} \vec{r}_{i+1/2} \quad (13)$$

becomes the nonlinear iteration

# Multiplicative Combination

The linear iteration

$$\vec{x}_{i+1/2} = \vec{x}_i - P^{-1} \vec{r}_i \quad (12)$$

$$\vec{x}_i = \vec{x}_{i+1/2} - Q^{-1} \vec{r}_{i+1/2} \quad (13)$$

becomes the nonlinear iteration

$$\vec{x}_{i+1} = \mathcal{M}(\mathcal{F}, \mathcal{N}(\mathcal{F}, \vec{x}_i, \vec{b}), \vec{b}) \quad (14)$$

# Nonlinear Right Preconditioning

For the linear case, we have

$$AP^{-1}\vec{y} = \vec{b} \quad (15)$$

$$\vec{x} = P^{-1}\vec{y} \quad (16)$$

so we define the preconditioning operation as

$$\vec{y} = \mathcal{M}(\mathcal{F}(\mathcal{N}(\mathcal{F}, \cdot, \vec{b})), \vec{x}_i, \vec{b}) \quad (17)$$

$$\vec{x} = \mathcal{N}(\mathcal{F}, \vec{y}, \vec{b}) \quad (18)$$

# Nonlinear Preconditioning

Type	Sym	Statement	Abbreviation
Additive	+	$\vec{x} + \alpha(\mathcal{M}(\mathcal{F}, \vec{x}, \vec{b}) - \vec{x}) + \beta(\mathcal{N}(\mathcal{F}, \vec{x}, \vec{b}) - \vec{x})$	$\mathcal{M} + \mathcal{N}$
Multiplicative	*	$\mathcal{M}(\mathcal{F}, \mathcal{N}(\mathcal{F}, \vec{x}, \vec{b}), \vec{b})$	$\mathcal{M} * \mathcal{N}$
Left Prec.	$-_L$	$\mathcal{M}(\vec{x} - \mathcal{N}(\mathcal{F}, \vec{x}, \vec{b}), \vec{x}, \vec{b})$	$\mathcal{M} -_L \mathcal{N}$
Right Prec.	$-_R$	$\mathcal{M}(\mathcal{F}(\mathcal{N}(\mathcal{F}, \vec{x}, \vec{b})), \vec{x}, \vec{b})$	$\mathcal{M} -_R \mathcal{N}$
Inner Lin. Inv.	\	$\vec{y} = \vec{J}(\vec{x})^{-1} \vec{r}(\vec{x}) = \mathbf{K}(\vec{J}(\vec{x}), \vec{y}_0, \vec{b})$	$\mathcal{N} \backslash \mathbf{K}$

# Nonlinear Richardson

```
1: procedure NRICH( $\vec{F}$ ,  $\vec{x}_i$ ,  $\vec{b}$ )
2:    $\vec{d} = -\vec{r}(\vec{x}_i)$ 
3:    $\vec{x}_{i+1} = \vec{x}_i + \lambda \vec{d}$                                  $\triangleright \lambda$  determined by line search
4: end procedure
5: return  $\vec{x}_{i+1}$ 
```

# Line Search

Equivalent to  $\text{NRICH} -_L \mathcal{N}$ :

$\text{NRICH} -_L \mathcal{N}$

# Line Search

Equivalent to  $\text{NRICH} -_L \mathcal{N}$ :

$\text{NRICH} -_L \mathcal{N}$

$\text{NRICH}(\vec{x} - \mathcal{N}(\mathcal{F}, \vec{x}, \vec{b}), \vec{x}, \vec{b})$

# Line Search

Equivalent to  $\text{NRICH} -_L \mathcal{N}$ :

$\text{NRICH} -_L \mathcal{N}$

$\text{NRICH}(\vec{x} - \mathcal{N}(\mathcal{F}, \vec{x}, \vec{b}), \vec{x}, \vec{b})$

$$\vec{x}_{i+1} = \vec{x}_i - \lambda \vec{r}_L$$

# Line Search

Equivalent to  $\text{NRICH} -_L \mathcal{N}$ :

$\text{NRICH} -_L \mathcal{N}$

$\text{NRICH}(\vec{x} - \mathcal{N}(\mathcal{F}, \vec{x}, \vec{b}), \vec{x}, \vec{b})$

$$\vec{x}_{i+1} = \vec{x}_i - \lambda \vec{r}_L$$

$$\vec{x}_{i+1} = \vec{x}_i + \lambda(\mathcal{N}(\mathcal{F}, \vec{x}_i, \vec{b}) - \vec{x}_i)$$

# PETSc Line Search

## BT Standard cubic back-tracking

- Defaults to full step when Wolfe conditions satisfied
- No more work than necessary
- May stagnate
- Can be badly scaled apart from  $\mathcal{N}$

## L2 Secant minimization of residual

- Optimal damping in the residual direction
- Minimize  $\|\vec{r}(\vec{x} + \lambda\delta\vec{x})\|_2$

## CP Secant minimization of energy

- Appropriate when  $\mathcal{F}$  is the gradient of an energy function
- Looks for roots of  $\delta\vec{x}^T \mathcal{F}(\vec{x} + \lambda\delta\vec{x})$

# Nonlinear GMRES

```

1: procedure NGMRES( $\mathcal{F}$ ,  $\vec{x}_i \cdots \vec{x}_{i-m+1}$ ,  $\vec{b}$ )
2:    $\vec{d}_i = -\vec{r}(\vec{x}_i)$ 
3:    $\vec{x}_i^M = \vec{x}_i + \lambda \vec{d}_i$ 
4:    $\mathcal{F}_i^M = \vec{r}(\vec{x}_i^M)$ 
5:   minimize  $\|\vec{r}((1 - \sum_{k=i-m}^{i-1} \alpha_k) \vec{x}_i^M + \sum_{k=i-m}^{i-1} \alpha_k \vec{x}_k)\|_2$  over
      $\{\alpha_{i-m} \cdots \alpha_{i-1}\}$ 
6:    $\vec{x}_i^A = (1 - \sum_{k=i-m}^{i-1} \alpha_k) \vec{x}_i^M + \sum_{k=i-m}^{i-1} \alpha_k \vec{x}_k$ 
7:    $\vec{x}_{i+1} = \vec{x}_i^A$  or  $\vec{x}_i^M$  if  $\vec{x}_i^A$  is insufficient.
8: end procedure
9: return  $\vec{x}_{i+1}$ 

```

Can emulate Anderson mixing and DIIS

# Nonlinear GMRES

```

1: procedure NGMRES( $\mathcal{F}$ ,  $\vec{x}_i \cdots \vec{x}_{i-m+1}$ ,  $\vec{b}$ )
2:    $\vec{d}_i = -\vec{r}(\vec{x}_i)$ 
3:    $\vec{x}_i^M = \vec{x}_i + \lambda \vec{d}_i$ 
4:    $\mathcal{F}_i^M = \vec{r}(\vec{x}_i^M)$ 
5:   minimize  $\|\vec{r}((1 - \sum_{k=i-m}^{i-1} \alpha_k) \vec{x}_i^M + \sum_{k=i-m}^{i-1} \alpha_k \vec{x}_k)\|_2$  over
      $\{\alpha_{i-m} \cdots \alpha_{i-1}\}$ 
6:    $\vec{x}_i^A = (1 - \sum_{k=i-m}^{i-1} \alpha_k) \vec{x}_i^M + \sum_{k=i-m}^{i-1} \alpha_k \vec{x}_k$ 
7:    $\vec{x}_{i+1} = \vec{x}_i^A$  or  $\vec{x}_i^M$  if  $\vec{x}_i^A$  is insufficient.
8: end procedure
9: return  $\vec{x}_{i+1}$ 

```

Can emulate Anderson mixing and DIIS

# Newton-Krylov

```
1: procedure  $\mathcal{N}\backslash K(\vec{F}, \vec{x}_i, \vec{b})$ 
2:    $\vec{d} = \vec{J}(\vec{x}_i)^{-1} \vec{r}(\vec{x}_i, \vec{b})$             $\triangleright$  solve by Krylov method
3:    $\vec{x}_{i+1} = \vec{x}_i + \lambda \vec{d}$             $\triangleright \lambda$  determined by line search
4: end procedure
5: return  $\vec{x}_{i+1}$ 
```

# Left Preconditioned Newton-Krylov

```
1: procedure  $\mathcal{N}\backslash K(\vec{x} - \vec{M}(\mathcal{F}, \vec{x}, \vec{b}), \vec{x}_i, 0)$ 
2:    $\vec{d} = \frac{\partial(\vec{x}_i - \mathcal{M}(\mathcal{F}, \vec{x}_i, \vec{b}))}{\partial \vec{x}_i}^{-1} (\vec{x}_i - \mathcal{M}(\mathcal{F}, \vec{x}_i, \vec{b}))$ 
3:    $\vec{x}_{i+1} = \vec{x}_i + \lambda \vec{d}$ 
4: end procedure
5: return  $\vec{x}_{i+1}$ 
```

# Jacobian Computation

$$\frac{\partial(\vec{x} - \mathcal{M}(\mathcal{F}, \vec{x}_i, \vec{b}))}{\vec{x}_i} = I - \frac{\partial \mathcal{M}(\mathcal{F}, \vec{x}_i, \vec{b})}{\partial \vec{x}_i},$$

Direct differencing would require

- one inner nonlinear iteration per **Krylov** iteration.

# Jacobian Computation

$$\frac{\partial(\vec{x} - \mathcal{M}(\mathcal{F}, \vec{x}_i, \vec{b}))}{\vec{x}_i} = I - \frac{\partial \mathcal{M}(\mathcal{F}, \vec{x}_i, \vec{b})}{\partial \vec{x}_i},$$

Direct differencing would require

- one inner nonlinear iteration per **Krylov** iteration.

# Jacobian Computation

**Impractical!**

$$\frac{\partial(\vec{x} - \mathcal{M}(\mathcal{F}, \vec{x}_i, \vec{b}))}{\vec{x}_i} = I - \frac{\partial \mathcal{M}(\mathcal{F}, \vec{x}_i, \vec{b})}{\partial \vec{x}_i},$$

Direct differencing would require

- one inner nonlinear iteration per **Krylov** iteration.

# Jacobian Computation

Approximation for NASM

$$\begin{aligned}\frac{\partial(\vec{x} - \mathcal{M}(\mathcal{F}, \vec{x}, \vec{b}))}{\partial \vec{x}} &= \frac{\partial(\vec{x} - (\vec{x} - \sum_b J_b(\vec{x}_b)^{-1} \mathcal{F}_b(\vec{x}_b)))}{\partial \vec{x}} \\ &\approx \sum_b J_b(\vec{x}_{b*})^{-1} J(\vec{x})\end{aligned}$$

This would require

- one inner nonlinear iteration
- small number of block solves

per **outer nonlinear** iteration.

X.-C. Cai and D. E. Keyes, *SIAM J. Sci. Comput.*, 24 (2002), pp. 183–200

# Right Preconditioned Newton-Krylov

```
1: procedure NK( $\vec{F}(\vec{M}(\vec{F}, \cdot, \vec{b}))$ ,  $\vec{y}_i, \vec{b}$ )
2:    $\vec{x}_i = \vec{M}(\vec{F}, \vec{y}_i, \vec{b})$ 
3:    $\vec{d} = \vec{J}(\vec{x})^{-1} \vec{r}(\vec{x}_i)$ 
4:    $\vec{x}_{i+1} = \vec{x}_i + \lambda \vec{d}$                                  $\triangleright \lambda$  determined by line search
5: end procedure
6: return  $\vec{x}_{i+1}$ 
```

# Jacobian Computation

## First-Order Approximation

Only the action of the original Jacobian is needed at first order:

$$\vec{y}_{i+1} = \vec{y}_i - \lambda \frac{\partial \mathcal{M}(\mathcal{F}, \vec{y}_i)}{\partial \vec{y}_i}^{-1} J(\mathcal{M}(\mathcal{F}, \vec{y}_i))^{-1} \mathcal{F}(\mathcal{M}(\mathcal{F}, \vec{y}_i))$$

$$\begin{aligned}\mathcal{M}(\mathcal{F}, \vec{y}_{i+1}) &= \mathcal{M}(\mathcal{F}, \vec{y}_i) - \lambda \frac{\partial \mathcal{M}(\mathcal{F}, \vec{y}_i)}{\partial \vec{y}_i}^{-1} J(\mathcal{M}(\mathcal{F}, \vec{y}_i))^{-1} \mathcal{F}(\mathcal{M}(\mathcal{F}, \vec{y}_i)) \\ &\approx \mathcal{M}(\mathcal{F}, \vec{y}_i)\end{aligned}$$

$$\begin{aligned}&- \lambda \frac{\partial \mathcal{M}(\mathcal{F}, \vec{y}_i)}{\partial \vec{y}_i} \frac{\partial \mathcal{M}(\mathcal{F}, \vec{y}_i)}{\partial \vec{y}_i}^{-1} J(\mathcal{M}(\mathcal{F}, \vec{y}_i))^{-1} \mathcal{F}(\mathcal{M}(\mathcal{F}, \vec{y}_i)) \\ &= \mathcal{M}(\mathcal{F}, \vec{y}_i) - \lambda J(\mathcal{M}(\mathcal{F}, \vec{y}_i))^{-1} \mathcal{F}(\mathcal{M}(\mathcal{F}, \vec{y}_i))\end{aligned}$$

$$\vec{x}_{i+1} = \vec{x}_i - \lambda J(\vec{x}_i)^{-1} \mathcal{F}(\vec{x}_i)$$

$\mathcal{N} \setminus K -_R \vec{M}$  is equivalent to  $\mathcal{N} \setminus K * \vec{M}$  at first order

# Jacobian Computation

## First-Order Approximation

Only the action of the original Jacobian is needed at first order:

$$\vec{y}_{i+1} = \vec{y}_i - \lambda \frac{\partial \mathcal{M}(\mathcal{F}, \vec{y}_i)}{\partial \vec{y}_i}^{-1} J(\mathcal{M}(\mathcal{F}, \vec{y}_i))^{-1} \mathcal{F}(\mathcal{M}(\mathcal{F}, \vec{y}_i))$$

$$\begin{aligned}\mathcal{M}(\mathcal{F}, \vec{y}_{i+1}) &= \mathcal{M}(\mathcal{F}, \vec{y}_i) - \lambda \frac{\partial \mathcal{M}(\mathcal{F}, \vec{y}_i)}{\partial \vec{y}_i}^{-1} J(\mathcal{M}(\mathcal{F}, \vec{y}_i))^{-1} \mathcal{F}(\mathcal{M}(\mathcal{F}, \vec{y}_i)) \\ &\approx \mathcal{M}(\mathcal{F}, \vec{y}_i)\end{aligned}$$

$$\begin{aligned}&- \lambda \frac{\partial \mathcal{M}(\mathcal{F}, \vec{y}_i)}{\partial \vec{y}_i} \frac{\partial \mathcal{M}(\mathcal{F}, \vec{y}_i)}{\partial \vec{y}_i}^{-1} J(\mathcal{M}(\mathcal{F}, \vec{y}_i))^{-1} \mathcal{F}(\mathcal{M}(\mathcal{F}, \vec{y}_i)) \\ &= \mathcal{M}(\mathcal{F}, \vec{y}_i) - \lambda J(\mathcal{M}(\mathcal{F}, \vec{y}_i))^{-1} \mathcal{F}(\mathcal{M}(\mathcal{F}, \vec{y}_i))\end{aligned}$$

$$\vec{x}_{i+1} = \vec{x}_i - \lambda J(\vec{x}_i)^{-1} \mathcal{F}(\vec{x}_i)$$

$\mathcal{N} \setminus K -_R \vec{M}$  is equivalent to  $\mathcal{N} \setminus K * \vec{M}$  at first order

# Jacobian Computation

## Direct Approximation

$$\begin{aligned}\mathcal{F}(\mathcal{M}(\mathcal{F}, \vec{y}_i, \vec{b})) &= J(\mathcal{M}(\mathcal{F}, \vec{y}_i, \vec{b})) \frac{\partial \mathcal{M}(\mathcal{F}, \vec{y}_i, \vec{b})}{\partial \vec{y}_i} (\vec{y}_{i+1} - \vec{y}_i) \\ &\approx J(\mathcal{M}(\mathcal{F}, \vec{y}_i, \vec{b})) (\mathcal{M}(\mathcal{F}, \vec{y}_i + \vec{d}, \vec{b}) - \vec{x}_i)\end{aligned}$$

- Solve for  $\vec{d}$
- Requires inner nonlinear solve for each Krylov iterate
- Needs FGMRES

P. Birken and A. Jameson, *J. Num. Meth. in Fluids*, 62 (2010), pp. 565–573

# Full Approximation Scheme (FAS)

## Nonlinear Multigrid

```
1: procedure FAS( $\vec{F}, \vec{x}_i, \vec{b}$ )
2:    $\vec{x}_s = \mathcal{M}_s(\mathcal{F}, \vec{x}_i, \vec{b})$ 
3:    $\vec{x}_c = \widehat{\mathbf{R}}\vec{x}_s$ 
4:    $\vec{b}_c = \mathcal{F}_c(\vec{x}_c) - \mathbf{R}[\mathcal{F}(\vec{x}_s) - \vec{b}]$ 
5:    $\vec{x}_s = \vec{x}_s + \mathbf{P}[\text{FAS}(\vec{F}_c, \vec{x}_c, \vec{b}_c) - \vec{x}_c]$ 
6:    $\vec{x}_{i+1} = \mathcal{M}_s(\mathcal{F}, \vec{x}_s, \vec{b})$ 
7: end procedure
8: return  $\vec{x}_{i+1}$ 
```

# Other Nonlinear Solvers

**NASM** Nonlinear Additive Schwarz

**NGS** Nonlinear Gauss-Siedel

**NCG** Nonlinear Conjugate Gradients

**QN** Quasi-Newton methods

# Outline

## 1 Algorithms

## 2 Experiments

- Composition
- Multilevel

# Outline

2

## Experiments

- Composition
- Multilevel

# SNES ex16

## 3D Large Deformation Elasticity

$$\int_{\Omega} \mathbf{F} \cdot \mathbf{S} : \nabla \mathbf{v} d\Omega + \int_{\Omega} \text{loading } \mathbf{e}_y \cdot \mathbf{v} d\Omega = 0 \quad (19)$$

$\mathbf{F}$  Deformation gradient

$\mathbf{S}$  Second Piola-Kirchhoff tensor

Saint Venant-Kirchhoff model of hyperelasticity


$\Omega$  -arc *angle* subsection of a cylindrical shell

-height *thickness*

-rad *inner radius*


-width *width*

# Large Deformation Elasticity




Unstressed and stressed configurations for the elasticity test problem.

# Large Deformation Elasticity



Coloration indicates vertical displacement in meters.

# Large Deformation Elasticity



P. Wriggers, Nonlinear Finite Element Methods, Springer, 2008.

# Large Deformation Elasticity


Running

## SNES example 16:

```
cd src/snes/examples/tutorials  
make ex16  
. /ex16 -da_grid_x 401 -da_grid_y 9 -da_grid_z 9  
-height 3 -width 3  
-rad 100 -young 100 -poisson 0.2  
-loading -1 -ploading 0
```

# Plain SNES Convergence

( $\mathcal{N} \setminus K - MG$ ) and NCG




# Plain SNES Convergence

Solver	T	N. It	L. It	Func	Jac	PC	NPC
NCG	53.05	4495	0	8991	—	—	—
( $\mathcal{N}\backslash K - MG$ )	23.43	27	1556	91	27	1618	—

# Composed SNES Convergence

NCG(10) + ( $\mathcal{N}\backslash K - MG$ ) and NCG(10) \* ( $\mathcal{N}\backslash K - MG$ )




# Composed SNES Convergence

Solver	T	N. It	L. It	Func	Jac	PC	NPC
NCG	53.05	4495	0	8991	—	—	—
( $\mathcal{N} \backslash K - MG$ )	23.43	27	1556	91	27	1618	—
NCG(10) + ( $\mathcal{N} \backslash K - MG$ )	14.92	9	459	218	9	479	—
NCG(10) * ( $\mathcal{N} \backslash K - MG$ )	16.34	11	458	251	11	477	—

# Preconditioned SNES Convergence

NGMRES- $R$  ( $\mathcal{N} \setminus K - MG$ ) and NCG- $L$  ( $\mathcal{N} \setminus K - MG$ )



# Preconditioned SNES Convergence

Solver	T	N. It	L. It	Func	Jac	PC	NPC
NCG	53.05	4495	0	8991	—	—	—
$(\mathcal{N}\backslash K - MG)$	23.43	27	1556	91	27	1618	—
NCG(10)	14.92	9	459	218	9	479	—
$+(\mathcal{N}\backslash K - MG)$							
NCG(10)	16.34	11	458	251	11	477	—
$\ast(\mathcal{N}\backslash K - MG)$							
NGMRES	9.65	13	523	53	13	548	13
$-R(\mathcal{N}\backslash K - MG)$							
NCG	9.84	13	529	53	13	554	13
$-L(\mathcal{N}\backslash K - MG)$							

# Outline

2

## Experiments

- Composition
- Multilevel

## SNES ex19

Driven Cavity Flow

$$-\Delta \vec{u} + \nabla \times \Omega = 0$$

$$-\Delta \Omega + \nabla \cdot (\vec{u} \Omega) - GR \nabla_x T = 0$$

$$-\Delta T + PR \nabla \cdot (\vec{u} T) = 0$$

# SNES ex19

## Driven Cavity Flow



$$-\Delta \vec{u} + \nabla \times \Omega = 0$$

$$\nabla \cdot (\vec{u}\Omega) - GR\nabla_x T = 0$$

$$-\Delta T + PR\nabla \cdot (\vec{u}T) = 0$$

# Driven Cavity Problem

## SNES ex19.c

```
./ex19 -lidvelocity 100 -grashof 1e2  
-da_grid_x 16 -da_grid_y 16 -da_refine 2  
-snes_monitor_short -snes_converged_reason -snes_view
```

# Driven Cavity Problem

## SNES ex19.c

```
./ex19 -lidvelocity 100 -grashof 1e2  
-da_grid_x 16 -da_grid_y 16 -da_refine 2  
-snes_monitor_short -snes_converged_reason -snes_view

lid velocity = 100, prandtl # = 1, grashof # = 100
0 SNES Function norm 768.116
1 SNES Function norm 658.288
2 SNES Function norm 529.404
3 SNES Function norm 377.51
4 SNES Function norm 304.723
5 SNES Function norm 2.59998
6 SNES Function norm 0.00942733
7 SNES Function norm 5.20667e-08
Nonlinear solve converged due to CONVERGED_FNORM_RELATIVE iterations 7
```

# Driven Cavity Problem

## SNES ex19.c

```
./ex19 -lidvelocity 100 -grashof 1e4  
-da_grid_x 16 -da_grid_y 16 -da_refine 2  
-snes_monitor_short -snes_converged_reason -snes_view
```

# Driven Cavity Problem

## SNES ex19.c

```
./ex19 -lidvelocity 100 -grashof 1e4  
-da_grid_x 16 -da_grid_y 16 -da_refine 2  
-snes_monitor_short -snes_converged_reason -snes_view

lid velocity = 100, prandtl # = 1, grashof # = 10000
0 SNES Function norm 785.404
1 SNES Function norm 663.055
2 SNES Function norm 519.583
3 SNES Function norm 360.87
4 SNES Function norm 245.893
5 SNES Function norm 1.8117
6 SNES Function norm 0.00468828
7 SNES Function norm 4.417e-08
Nonlinear solve converged due to CONVERGED_FNORM_RELATIVE iterations 7
```

# Driven Cavity Problem

## SNES ex19.c

```
./ex19 -lidvelocity 100 -grashof 1e5  
-da_grid_x 16 -da_grid_y 16 -da_refine 2  
-snes_monitor_short -snes_converged_reason -snes_view
```

# Driven Cavity Problem

## SNES ex19.c

```
./ex19 -lidvelocity 100 -grashof 1e5  
-da_grid_x 16 -da_grid_y 16 -da_refine 2  
-snes_monitor_short -snes_converged_reason -snes_view  
  
lid velocity = 100, prandtl # = 1, grashof # = 100000  
0 SNES Function norm 1809.96  
Nonlinear solve did not converge due to DIVERGED_LINEAR_SOLVE iterations 0
```

# Driven Cavity Problem

## SNES ex19.c

```
./ex19 -lidvelocity 100 -grashof 1e5  
-da_grid_x 16 -da_grid_y 16 -da_refine 2 -pc_type lu  
-snes_monitor_short -snes_converged_reason -snes_view
```

```
lid velocity = 100, prandtl # = 1, grashof # = 100000  
0 SNES Function norm 1809.96  
1 SNES Function norm 1678.37  
2 SNES Function norm 1643.76  
3 SNES Function norm 1559.34  
4 SNES Function norm 1557.6  
5 SNES Function norm 1510.71  
6 SNES Function norm 1500.47  
7 SNES Function norm 1498.93  
8 SNES Function norm 1498.44  
9 SNES Function norm 1498.27  
10 SNES Function norm 1498.18  
11 SNES Function norm 1498.12  
12 SNES Function norm 1498.11  
13 SNES Function norm 1498.11  
14 SNES Function norm 1498.11  
...
```

# Nonlinear Preconditioning

```
./ex19 -lidvelocity 100 -grashof 5e4 -da_refine 4 -snes_monitor_short  
-snes_type newtonls -snes_converged_reason  
-pc_type lu
```

```
lid velocity = 100, prandtl # = 1, grashof # = 50000
0 SNES Function norm 1228.95
1 SNES Function norm 1132.29
2 SNES Function norm 1026.17
3 SNES Function norm 925.717
4 SNES Function norm 924.778
5 SNES Function norm 836.867
...
21 SNES Function norm 585.143
22 SNES Function norm 585.142
23 SNES Function norm 585.142
24 SNES Function norm 585.142
...
```

# Nonlinear Preconditioning

```
./ex19 -lidvelocity 100 -grashof 5e4 -da_refine 4 -snes_monitor_short  
-snes_type fas -snes_converged_reason  
-fas_levels_snes_type gs -fas_levels_snes_max_it 6
```

```
lid velocity = 100, prandtl # = 1, grashof # = 50000  
0 SNES Function norm 1228.95  
1 SNES Function norm 574.793  
2 SNES Function norm 513.02  
3 SNES Function norm 216.721  
4 SNES Function norm 85.949
```

```
Nonlinear solve did not converge due to DIVERGED_INNER iterations 4
```

# Nonlinear Preconditioning

```
./ex19 -lidvelocity 100 -grashof 5e4 -da_refine 4 -snes_monitor_short  
-snes_type fas -snes_converged_reason  
-fas_levels_snes_type gs -fas_levels_snes_max_it 6  
-fas_coarse_snes_converged_reason
```

```
lid velocity = 100, prandtl # = 1, grashof # = 50000  
0 SNES Function norm 1228.95  
    Nonlinear solve converged due to CONVERGED_FNORM_RELATIVE its 12  
1 SNES Function norm 574.793  
    Nonlinear solve did not converge due to DIVERGED_MAX_IT its 50  
2 SNES Function norm 513.02  
    Nonlinear solve did not converge due to DIVERGED_MAX_IT its 50  
3 SNES Function norm 216.721  
    Nonlinear solve converged due to CONVERGED_FNORM_RELATIVE its 22  
4 SNES Function norm 85.949  
    Nonlinear solve did not converge due to DIVERGED_LINE_SEARCH its 42  
Nonlinear solve did not converge due to DIVERGED_INNER iterations 4
```

# Nonlinear Preconditioning

```
./ex19 -lidvelocity 100 -grashof 5e4 -da_refine 4 -snes_monitor_short  
-snes_type fas -snes_converged_reason  
-fas_levels_snes_type gs -fas_levels_snes_max_it 6  
-fas_coarse_snes_linesearch_type basic  
-fas_coarse_snes_converged_reason
```

```
lid velocity = 100, prandtl # = 1, grashof # = 50000  
0 SNES Function norm 1228.95  
    Nonlinear solve converged due to CONVERGED_FNORM_RELATIVE its 6  
:  
47 SNES Function norm 78.8401  
    Nonlinear solve converged due to CONVERGED_FNORM_RELATIVE its 5  
48 SNES Function norm 73.1185  
    Nonlinear solve converged due to CONVERGED_FNORM_RELATIVE its 6  
49 SNES Function norm 78.834  
    Nonlinear solve converged due to CONVERGED_FNORM_RELATIVE its 5  
50 SNES Function norm 73.1176  
    Nonlinear solve converged due to CONVERGED_FNORM_RELATIVE its 6  
:
```

# Nonlinear Preconditioning

```
./ex19 -lidvelocity 100 -grashof 5e4 -da_refine 4 -snes_monitor_short  
-snes_type nrichardson -npc_snes_max_it 1 -snes_converged_reason  
-npc_snes_type fas -npc_fas_coarse_snes_converged_reason  
-npc_fas_levels_snes_type gs -npc_fas_levels_snes_max_it 6  
-npc_fas_coarse_snes_linesearch_type basic
```

```
lid velocity = 100, prandtl # = 1, grashof # = 50000  
0 SNES Function norm 1228.95  
    Nonlinear solve converged due to CONVERGED_FNORM_RELATIVE its 6  
1 SNES Function norm 552.271  
    Nonlinear solve converged due to CONVERGED_FNORM_RELATIVE its 27  
2 SNES Function norm 173.45  
    Nonlinear solve converged due to CONVERGED_FNORM_RELATIVE its 45  
:  
43 SNES Function norm 3.45407e-05  
    Nonlinear solve converged due to CONVERGED_SNORM_RELATIVE its 2  
44 SNES Function norm 1.6141e-05  
    Nonlinear solve converged due to CONVERGED_SNORM_RELATIVE its 2  
45 SNES Function norm 9.13386e-06  
Nonlinear solve converged due to CONVERGED_FNORM_RELATIVE iterations 45
```



# Nonlinear Preconditioning

```
./ex19 -lidvelocity 100 -grashof 5e4 -da_refine 4 -snes_monitor_short  
-snes_type ngmres -npc_snes_max_it 1 -snes_converged_reason  
-npc_snes_type fas -npc_fas_coarse_snes_converged_reason  
-npc_fas_levels_snes_type gs -npc_fas_levels_snes_max_it 6  
-npc_fas_coarse_snes_linesearch_type basic
```

```
lid velocity = 100, prandtl # = 1, grashof # = 50000  
0 SNES Function norm 1228.95  
    Nonlinear solve converged due to CONVERGED_FNORM_RELATIVE its 6  
1 SNES Function norm 538.605  
    Nonlinear solve converged due to CONVERGED_FNORM_RELATIVE its 13  
2 SNES Function norm 178.005  
    Nonlinear solve converged due to CONVERGED_FNORM_RELATIVE its 24  
:  
27 SNES Function norm 0.000102487  
    Nonlinear solve converged due to CONVERGED_FNORM_RELATIVE its 2  
28 SNES Function norm 4.2744e-05  
    Nonlinear solve converged due to CONVERGED_SNORM_RELATIVE its 2  
29 SNES Function norm 1.01621e-05  
Nonlinear solve converged due to CONVERGED_FNORM_RELATIVE iterations 29
```



# Nonlinear Preconditioning

```
./ex19 -lidvelocity 100 -grashof 5e4 -da_refine 4 -snes_monitor_short  
-snes_type ngmres -npc_snes_max_it 1 -snes_converged_reason  
-npc_snes_type fas -npc_fas_coarse_snes_converged_reason  
-npc_fas_levels_snes_type newtonls -npc_fas_levels_snes_max_it 6  
-npc_fas_levels_snes_linesearch_type basic  
-npc_fas_levels_snes_max_linear_solve_fail 30  
-npc_fas_levels_ksp_max_it 20 -npc_fas_levels_snes_converged_reason  
-npc_fas_coarse_snes_linesearch_type basic  
lid velocity = 100, prandtl # = 1, grashof # = 50000  
0 SNES Function norm 1228.95  
    Nonlinear solve did not converge due to DIVERGED_MAX_IT its 6  
:  
    Nonlinear solve converged due to CONVERGED_SNORM_RELATIVE its 1  
:  
1 SNES Function norm 0.1935  
2 SNES Function norm 0.0179938  
3 SNES Function norm 0.00223698  
4 SNES Function norm 0.000190461  
5 SNES Function norm 1.6946e-06  
Nonlinear solve converged due to CONVERGED_FNORM_RELATIVE iterations 5
```

# Nonlinear Preconditioning

```
./ex19 -lidvelocity 100 -grashof 5e4 -da_refine 4 -snes_monitor_short  
-snes_type composite -snes_composite_type additiveoptimal  
-snes_composite_sneses fas,newtonls -snes_converged_reason  
-sub_0_fas_levels_snes_type gs -sub_0_fas_levels_snes_max_it 6  
-sub_0_fas_coarse_snes_linesearch_type basic  
-sub_1_snes_linesearch_type basic -sub_1_pc_type mg
```

```
lid velocity = 100, prandtl # = 1, grashof # = 50000
```

```
0 SNES Function norm 1228.95  
1 SNES Function norm 541.462  
2 SNES Function norm 162.92  
3 SNES Function norm 48.8138  
4 SNES Function norm 11.1822  
5 SNES Function norm 0.181469  
6 SNES Function norm 0.00170909  
7 SNES Function norm 3.24991e-08
```

```
Nonlinear solve converged due to CONVERGED_FNORM_RELATIVE iterations 7
```

# Nonlinear Preconditioning

```
./ex19 -lidvelocity 100 -grashof 5e4 -da_refine 4 -snes_monitor_short  
-snes_type composite -snes_composite_type multiplicative  
-snes_composite_sneses fas,newtonls -snes_converged_reason  
-sub_0_fas_levels_snes_type gs -sub_0_fas_levels_snes_max_it 6  
-sub_0_fas_coarse_snes_linesearch_type basic  
-sub_1_snes_linesearch_type basic -sub_1_pc_type mg
```

```
lid velocity = 100, prandtl # = 1, grashof # = 50000
```

```
0 SNES Function norm 1228.95  
1 SNES Function norm 544.404  
2 SNES Function norm 18.2513  
3 SNES Function norm 0.488689  
4 SNES Function norm 0.000108712  
5 SNES Function norm 5.68497e-08
```

```
Nonlinear solve converged due to CONVERGED_FNORM_RELATIVE iterations 5
```

# Nonlinear Preconditioning

Solver	T	N. It	L. It	Func	Jac	PC	NPC
$(\mathcal{N}\backslash K - MG)$	9.83	17	352	34	85	370	–
$NGMRES - R$	7.48	10	220	21	50	231	10
$(\mathcal{N}\backslash K - MG)$							
FAS	6.23	162	0	2382	377	754	–
FAS + $(\mathcal{N}\backslash K - MG)$	8.07	10	197	232	90	288	–
FAS * $(\mathcal{N}\backslash K - MG)$	4.01	5	80	103	45	125	–
NRICH - L FAS	3.20	50	0	1180	192	384	50
NGMRES - R FAS	1.91	24	0	447	83	166	24

# Nonlinear Preconditioning

See discussion in:

## **Composing scalable nonlinear solvers,**

Peter Brune, Matthew Knepley, Barry Smith, and Xuemin Tu,

ANL/MCS-P2010-0112, Argonne National Laboratory, 2012.

<http://www.mcs.anl.gov/uploads/cels/papers/P2010-0112.pdf>